




Opioid Addiction


R. Corey Waller MD, MS









Opioid Addiction


R. Corey Waller MD, MS



Disease of Addiction







Behavior

Diagnosis based in
the description of
behavior

Aberrant behavior
should be
expected

Therefore behavior
is a symptom not a
frustration



Heroin Use

Smoked



Dosing

1 dime bag = about 0.1 gram and generally equals 10-20 mg of morphine

10 dime bags = 1 gram

They will use 3 to 8 times per day depending on purity, tolerance and what it is cut with.

Most people will use 1/2 to 1 gram a day

Injected



Injected



Smoked



Dosing

1 dime bag = about 0.1 gram and generally equals 10-20 mg of morphine

10 dime bags = 1 gram

They will use 3 to 8 times per day depending on purity, tolerance and what it is cut with.

Most people will use 1/2 to 1 gram a day

How do we treat it?

Medication Assisted Treatment

- Methadone
- Buprenorphine
- Naltrexone (Oral or 30 day injection)

Behavioral Therapies

- Cognitive Behavioral Therapy
- Contingency Management
- Motivational Enhancement Therapy
- Treat co-morbid conditions

12-step?

Medication Assisted Treatment

- Methadone
- Buprenorphine
- Naltrexone (Oral or 30 day injection)

Behavioral Therapies

- Cognitive Behavioral Therapy
- Contingency Management
- Motivational Enhancement Therapy
- Treat co-morbid conditions

12-step?

Conclusions

New paradigm

Addiction is a chronic
Neurobiological disease

Many ways to use heroin

We **CAN** treat this
disease



Opioid Addiction

R. Corey Waller MD, MS

